## Dark -Field Microscopic Analysis on the Blood of 1,006 Symptomatic Persons After Anti-**COVID mRNA Injections from Pfizer/BioNtech** or Moderna

New large published study from Italy confirms extreme blood abnormalities and selfassembly nanostructures













Figure 4. This image at 120x magnification (3x magnification digitally produced) highlights a typical self-aggregating structuring in fibro/tubular mode.

This recently published study from Italy once again confirms self assembly nanostructures in the blood of C19 injected people. The authors correlate the blood findings with the symptoms of the affected. For example this case was discussed:

This individual is a male of 33 years, who formerly was an athlete, apparently healthy before inoculation with an mRNA Pfizer injection. One month after receiving the first dose of the Pfizer "vaccine", he showed marked asthenia, a constant gravitational headache (i.e., one sensitive to the position and movements of his head and body such that the pain was increased by movement of the head up or down). The headaches were unresponsive to common pain killers. Diffuse rheumatic arthralgia with dyspnea on exertion were noted.

This was his blood finding:




The authors show that these self-assembly nanoparticles conglomerate to create massive structures. These provide the basis for extensive clotting and abnormal blood flow.



Figure 6. The sharper and larger image on the left seems to suggest the power of the exogenous particles introduced into the blood of mRNA injection recipients to assemble themselves into massive structures; we can see in both (a) and (b) at 120x magnification evidence of what appear to be lamellar configurations similar to agglomerations occurring in a field of forces pulling colloidal particles in the plasma together. The relative bulk of the particle conglobate can be easily estimated by comparison with the crythrocytes at the periphery of the much larger mass. It was also measured accurately at 113.91µm by 139.99µm (see the hatch mark green lines in b) using the Delta-Pix Software as shown in the computer screen shot at the right.

Here are further self assembly nanostructures:




blood profile of 948 patients have never been observed after inoculation by any vaccines in the past according to our clinical experience.

The sudden transition, usually at the time of a second mRNA injection, from a state of perfect normalcy to a pathological one, with accompanying hemolysis, visible packing and stacking of red blood cells in conjunction with the formation of gigantic conglomerate foreign structures, some of them appearing as graphene-family super-structures, is unprecedented.


\*Such phenomena have never been seen before after any "vaccination" of the past.\*

In our collective experience, and in our shared professional opinion, the large quantity of particles in the blood of mRNA injection recipients is incompatible with normal blood flow especially at the level of the capillaries."

Here is the full article:



This is mounting evidence in addition to what already has been published with similar findings that I listed here:



In my view, the discovery of novel screening tools that can be used for early detection of potentially life threatening blood clots is imperative.

These are exactly the blood findings described by my colleague, Felipe Reitz. More than ever, I believe that the evaluation if Computerized Thermography Imaging could be an early screening tool is imperative. I wrote about this here, and please look at our video interview.



## Summary:

If these structures are magnetic, MRI's and other imaging studies should not be performed as they may cause tissue damage due to excitation of the magnetic particles. This could cause further harm and injury to the injected. Full body Ultrasound imaging is difficult to perform, but full body Computerized Thermography could be done safely.


Dangerously, many allopathic physicians treating vax injuries are in complete denial of the self assembly nanostructures as a cause for pathology. I wonder how much evidence they need to take this seriously?

We must find the safest way of screening for asymptomatic blood clotting, and clearly live blood analysis is a tool that all treating physicians should urgently familiarize themselves with. D-Dimers are valuable, but may not be able to pick up all blood clotting. Dr. Charles Hoffe stated that 62% of his C19 injected patients had an elevated D-Dimer, but all live blood analysis in injected patients are abnormal. The sensitivity of D-Dimer testing may be suboptimal.

Further correlation studies in C19 injected people with evaluation of live blood analysis, D-Dimer testing, Computerized Thermography Imaging and if abnormal, Doppler Ultrasound confirmation need to happen urgently. We need to screen every athlete, every pilot, every medical professional - and in fact every person who got theses shots. If we do, we may be able to prevent sudden adult or child death syndrome.

Above all, the dangerous shots need to be stopped immediately and everyone responsible, held accountable.





© 2023 Ana Mihalcea, MD, PhD · <u>Privacy</u> · <u>Terms</u> · <u>Collection notice</u>



Get the app